



| つ | |
|----------|--|
| ~ | |
| | |

| R | 1214 | 0 |
|---|------|---|
| | | |

Generic Name: acetaminophen and propoxyphene (a SEET a MIN oh fen and proe POX i feen)

Brand Name: Balacet, Darvocet A500, Darvocet-N 100, Darvocet-N 50, Propacet 100

Side Effects

Interactions

More ~

What is Propacet 100 (acetaminophen and propoxyphene)?

ADUERTISEMENT

Acetaminophen and propoxyphene was withdrawn from the U.S. market in November 2010.

Propoxyphene is in a group of drugs called narcotic pain relievers.

Acetaminophen is a less potent pain reliever and a fever reducer that increases the effects of propoxyphene.

The combination of acetaminophen and propoxyphene is used to relieve mild to moderate pain with or without fever.

Acetaminophen and propoxyphene may also be used for purposes not listed in thismedication of guide.

What is the most important information I should know about Propacet 100 (acetaminophen and propoxyphene)?

Acetaminophen and propoxyphene was withdrawn from the U.S. market in November 2010.

Propoxyphene may be habit-forming and **should be used only by the person it was prescribed for** Keep the medication in a secure place where others cannot get to it.



Do not use acetaminophen and propoxyphene if you have taken an MAO inhibitor such as furazolidone (Furoxone), isocarboxazid (Marplan), phenelzine (Nardil), rasagiline (Azilect), selegiline (Eldepryl, Emsam), or tranylcypromine (Parnate) in the last 14 days. Serious, life threatening side effects or can occur if you use acetaminophen and propoxyphene before the MAO inhibitor has cleared from your body or.

What should I discuss with my healthcare provider before taking Propacet 100 (acetaminophen and propoxyphene)?

Do not use this medication if you are allergic to acetaminophen (Tylenol) or propoxyphene.

Propoxyphene may be habit-forming and should be used only by the person it was prescribed for **Never share this medication** with another person, especially someone with a history of drug abuse or addiction. Keep the medication in a place where others cannot get to it.

Do not use acetaminophen and propoxyphene if you have used an MAO inhibitor such as furazolidone (Furoxone), isocarboxazid (Marplan), phenelzine (Nardil), rasagiline (Azilect), selegiline (Eldepryl, Emsam, Zelapar), or tranylcypromine (Parnate) in the last 14 days. A dangerous drug interaction could occur, leading to serious side effects.

To make sure you can safely take acetaminophen and propoxyphene, tell your doctor if you have any of these other conditions:

- asthma, COPD, sleep apnea, or other breathing disorders;
- · liver or kidney disease;
- a history of head injury or brain tumor;
- a gallbladder or pancreas disorder;
- a stomach or intestinal disorder;
- suicidal thoughts or behavior; or
- · mental illness, or a history of drug or alcohol addiction.

Tell your doctor if you drink more than three alcoholic beverages per day or if you have ever had alcoholic liver disease (cirrhosis). You may not be able to take medication that contains acetaminophen.

FDA pregnancy category C. It is not known whether this medication is harmful to an unborn<u>baby</u> $\[mathbb{Z}\]$, but it could cause breathing problems or addiction/withdrawal symptoms in a newborn. Tell your doctor if you are pregnant or plan to become pregnant during treatment.

Acetaminophen and propoxyphene can pass into breast milk and may harm a nursing baby. Do not use this medication without telling your doctor if you are breast-feeding a baby.

ADUERTISEMENT

How should I take Propacet 100 (acetaminophen and propoxyphene)?

Take exactly as prescribed by your doctor. Do not take in larger or smaller amounts or for longer than recommended. Follow the directions on your prescription label. Tell your doctor if the medicine seems to stop working as well in relieving your pain.

An overdose of acetaminophen can damage your liver. Adults should not take more than 1 gram (1000 mg) of acetaminophen per dose or 4 grams (4000 mg) per day. If you drink more than three alcoholic beverages per day, you should never use more than 2 grams (2000 mg) of acetaminophen per day.

Do not stop using acetaminophen and propoxyphene suddenly, or you could have unpleasant withdrawal symptoms. Ask your doctor how to avoid withdrawal symptoms when you stop using acetaminophen and propoxyphene.

If you need surgery, tell the surgeon ahead of time that you are using acetaminophen and propoxyphene. You may need to stop using the medicine for a short time.

Store at room temperature away from moisture and heat.

Keep track of the amount of medicine used from each new bottle. Propoxyphene is a drug of abuse and you should be aware if anyone is using your medicine improperly or without a prescription.

What happens if I miss a dose?

Since acetaminophen and propoxyphene is taken as needed, you may not be on a dosing schedule. If you are taking the medication regularly, take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222 An overdose of acetaminophen and propoxyphene can be fatal.

Overdose symptoms may include extreme drowsiness, nausea, vomiting, dark urine, jaundice (yellowing of the skin or eyes), pinpoint or dilated pupils, confusion, seizure (convulsions), cold and clammy skin, blue lips, weak pulse, slow or uneven heart rate, shallow breathing, fainting, or breathing that stops.

What should I avoid while taking Propacet 100 (acetaminophen and propoxyphene)?

This medication may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert.

Ask a doctor or pharmacist before using any other cold, allergy, pain, or sleep medication. Acetaminophen (sometimes abbreviated as "APAP") is contained in many combination medicines. Taking certain products together can cause you to get too much acetaminophen. Check the label to see if a medicine contains acetaminophen or APAP.

Drinking alcohol can increase certain side effects of acetaminophen and propoxyphene.

Propacet 100 (acetaminophen and propoxyphene) side effects

Get emergency medical help if you have any of these signs of an allergic reaction: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have a serious side effect such as:

- · shallow breathing, slow heartbeat, feeling like you might pass out;
- · chest pain, feeling short of breath;
- confusion, unusual thoughts or behavior;
- · seizure (convulsions); or
- nausea, stomach pain, loss of appetite, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes).

Less serious side effects include:

- · feeling dizzy or drowsy;
- mild nausea, vomiting, upset stomach, constipation;
- · headache, blurred vision; or
- dry mouth.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

See also: Side effects (in more detail)



What other drugs will affect Propacet 100 (acetaminophen and propoxyphene)?

Cold or allergy medicine, sleeping pills, muscle relaxers, antidepressants, or seizure medication can add to sleepiness caused by propoxyphene, or could slow your breathing. Tell your doctor if you regularly use any of these medicines, or any other narcotic medications.

Tell your doctor about all other medications you use, especially:

- aspirin or a blood thinner such as warfarin (Coumadin);
- · birth control pills;
- dexamethasone (Decadron, Hexadrol);
- a diuretic (water pill) such as furosemide (Lasix);
- St. John's wort:
- an antibiotic such as clarithromycin (Biaxin), erythromycin (E.E.S., EryPed, Ery-Tab, Erythrocin), rifabutin (Mycobutin), rifampin (Rifadin, Rifater, Rifamate), or rifapentine (Priftin);
- an antidepressant such as nefazodone;
- antifungal medication such as fluconazole (Diflucan), itraconazole (Sporanox), ketoconazole (Nizoral), or voriconazole (Vfend);
- a barbiturate such as phenobarbital (Solfoton);
- heart or blood pressure medication such as amiodarone (Cordarone, Pacerone), diltiazem (Cartia, Cardizem), nifedipine (Nifedical, Procardia), propranolol (Inderal), verapamil (Calan, Covera, Isoptin, Verelan), and others;
- HIV or AIDS medicine such as fosamprenavir (Lexiva), nelfinavir (Viracept), ritonavir (Norvir), zidovudine (Retrovir), and others: or
- seizure medication such as carbamazepine (Carbatrol, Tegretol), lamotrigine (Lamictal), phenytoin (Dilantin), and others.

This list is not complete and there are many other drugs that can interact with acetaminophen and propoxyphene. Tell your doctor about all the prescription and over-the-counter medications you use. This includes vitamins, minerals, herbal products, and drugs prescribed by other doctors. Do not start using a new medication without telling your doctor.














▲ Add to My Med List

More about Propacet 100 (acetaminophen / propoxyphene)

- Side Effects
- **During Pregnancy**
- Drug Images
- Drug Interactions
- Support Group
- 0 Reviews Add your own review/rating
- Consumer resources
- TINUED Other brands: Darvocet-N 100, Darvocet-N 50, Balacet, Darvocet A500, More (1) »
- Professional resources
- Propoxyphene and Acetaminophen Tablets (FDA)
- Related treatment guides
- Pain

Where can I get more information?

• Your pharmacist can provide more information about acetaminophen and propoxyphene.



Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Disclaimer: Every effort has been made to ensure that the information provided by Cerner Multum, Inc. ('Multum') is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum information has been compiled for use by healthcare practitioners and consumers in the United States and therefore Multum does not warrant that uses outside of the United States are appropriate, unless specifically indicated otherwise. Multum's drug information does not endorse drugs, diagnose patients or recommend therapy. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. Multum does not assume any responsibility for any aspect of healthcare administered with the aid of information Multum provides. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

Copyright 1996-2012 Cerner Multum, Inc. Version: 9.02 Revision Date: 2010-12-15, 5:01:39 PM.

DRUG STATUS



Availability Discontinued



Pregnancy Category Risk cannot be ruled out



CSA Schedule Some potential for abuse

| Manufacturer | ADVERTISEMENT |
|--|---------------------------|
| Teva Pharmaceuticals USA, Inc. | |
| | |
| Drug Class | |
| Narcotic analgesic combinations | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| Related Drugs | |
| Pain | |
| tramadol, oxycodone, acetaminophen, naproxen, Tylenol, aspirin, ibuprofen, amitriptyline, hydrocodone, Per | |
| hydroxyzine, diclofenac, Celebrex, fentanyl, morphine, Vicodin, OxyContin, Dilaudid, diphenhydramine topica More | al, methadone, Paracetamo |
| | |
| FD/A CONSUMER UPDATES | |
| FEAT CONSUMER OF DATES | |
| Fighting Allergy Season with Medications | |
| Medications for High Blood Pressure | |
| Irritable Bowel Syndrome Treatments Aren't One-Size-Fits-All | |
| Fighting Diabetes' Deadly Impact on Minorities | |
| More FDA updates | |
| | |
| Propacet 100 Rating | |
| No Reviews - Be the first! | 9 ./ 01 |

Propacet 100 Images



Propacet 100 (acetaminophen / propoxyphene) 650 mg / 100 mg



View larger images

Related News and Articles

Health Tip: Use Proper Lifting Technique

March 16, 2016

CDC Releases Guideline for Prescribing Opioids for Chronic Pain

March 15, 2016

Health Tip: Managing a Fever at Home

March 15, 2016

Related Questions & Answers

Does Propacet 100 keep you up at night (cause insomnia?)?

Read more questions... >





Drugs.com Mobile Apps

The easiest way to lookup drug information, identify pills, check interactions and set up your own personal medication records. Available for Android and iOS devices.

Explore Apps



SUPPORT

Help Center

Frequent Questions

Sitemap

Contact Us

ABOUT

About Drugs.com

Advertising Policy

Content Submissions

Drugs.com Blog

TERMS & PRIVACY

Editorial Policy

Privacy Policy

Terms of Use

Attribution & Citations

CONNECT











Subscribe to receive email notifications whenever new articles are published.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex® (updated Feb 24th, 2016), Cerner Multum™ (updated Mar 1st, 2016), Wolters Kluwer™ (updated Mar 1st, 2016) and others. To view content sources and attributions, please refer to our editorial policy.





Ad Choices



We comply with the HONcode standard for trustworthy health information - verify here