



Generic Name: clorazepate (klor AZ e pate) Brand Name: Tranxene SD, Tranxene T-Tab		
ROUERTISEMENT		
Overview		
Side Effects		
Dosage		
Interactions		
Professional		
More ✓		
What is Transons T Tab (clarazonats)?		

What is Tranxene T-Tab (clorazepate)?

ADUERTISEMENT

Clorazepate is in a group of drugs called benzodiazepines (ben-zoe-dye-AZE-eh-peens). Clorazepate affects chemicals in the brain that may become unbalanced and cause anxiety or seizures.

Clorazepate is used to treat anxiety disorders, partial seizures, or alcohol withdrawalsymptoms 丞.

# What is the most important information I should know about Tranxene T-Tab (clorazepate)?

Do not use this medication if you are allergic to clorazepate or to other benzodiazepines, such as alprazolam (Xanax), chlordiazepoxide (Librium), diazepam (Valium), lorazepam (Ativan), or oxazepam (Serax).

This medication can cause birth defects in an unborn baby. Do not use clorazepate if you are pregnant.



Before taking clorazepate, tell your doctor if you have any breathing problems, glaucoma, kidney or liver disease, or a history of depression, suicidal thoughts, or addiction to drugs or alcohol.

Do not drink alcohol while taking clorazepate. This medication can increase the effects of alcohol.

Avoid using other medicines that make you sleepy. They can add to sleepiness caused by clorazepate.

Clorazepate may be habit-forming and **should be used only by the person it was prescribed for** Clorazepate should never be shared with another person, especially someone who has a history of drug abuse or addiction. Keep the medication in a secure place where others cannot get to it.

# What should I discuss with my healthcare provider before taking Tranxene T-Tab (clorazepate)?

Do not use this medication if you have narrow-angle glaucoma, or if you are allergic to clorazepate or to other benzodiazepines, such as alprazolam (Xanax), chlordiazepoxide (Librium), diazepam (Valium), lorazepam (Ativan), or oxazepam (Serax).

Before taking clorazepate, tell your doctor if you are allergic to any drugs, or if you have:

- glaucoma;
- asthma, emphysema, bronchitis, chronic obstructive pulmonary disorder (COPD), or other breathing problems;
- kidney or liver disease;
- epilepsy or other seizure disorder;
- a history of depression or suicidal thoughts or behavior; or
- a history of drug or alcohol addiction.

If you have any of these conditions, you may need a dose adjustment or special tests during treatment.

Clorazepate can cause birth defects in an unborn baby. Do not use clorazepate without your doctor's consent if you are <a href="mailto:pregnant">pregnant</a> \overline{\mathbb{Z}}. Tell your doctor if you become pregnant during treatment. Use an effective form of birth control while you are using this medication.

Clorazepate can pass into breast milk and could harm a nursing baby. Do not use this medication without telling your doctor if you are breast-feeding a baby.

The sedative effects of clorazepate may last longer in older adults. Accidental falls are common in elderly patients who take benzodiazepines. Use caution to avoid falling or accidental injury while you are taking clorazepate.

Do not give this medication to a child younger than 9 years old.



# How should I take Tranxene T-Tab (clorazepate)?

Take this medication exactly as it was prescribed for you. Do not take the medication in larger amounts, or take it for longer than recommended by your doctor. Follow the directions on your prescription label.

Your doctor may occasionally change your dose to make sure you get the best results from this medication.

Clorazepate should be used for only a short time. Do not take this medication for longer than 4 months without your doctor's advice.

Clorazepate may be habit-forming and **should be used only by the person it was prescribed for** Clorazepate should never be shared with another person, especially someone who has a history of drug abuse or addiction. Keep the medication in a secure place where others cannot get to it.

Contact your doctor if this medicine seems to stop working as well in treating your anxiety symptoms.

Do not stop using clorazepate suddenly without first talking to your doctor. You may need to use less and less before you stop the medication completely.

Your symptoms may return when you stop using clorazepate after using it over a long period of time. You may also have withdrawal symptoms when you stop using clorazepate. Withdrawal symptoms may include tremor, sweating, trouble sleeping, muscle cramps, stomach pain, vomiting, diarrhea, irritability, memory problems, confusion, unusual thoughts or behavior, and seizure (convulsions).

To be sure this medication is not causing harmful effects, your blood and liver function may need to be tested on a regular basis. Do not miss any scheduled visits to your doctor.

Store clorazepate at room temperature away from moisture, heat, and light.

Keep track of how many pills have been used from each new bottle of this medicine. Benzodiazepines are drugs of abuse and you should be aware if any person in the household is using this medicine improperly or without a prescription.

# What happens if I miss a dose?

Take the missed dose as soon as you remember. If it is almost time for your next dose, skip the missed dose and take the medicine at your next regularly scheduled time. Do not take extra medicine to make up the missed dose.

# What happens if I overdose?

Seek emergency medical attention if you think you have used too much of this medicine **An overdose of clorazepate can be fatal.** 

Overdose symptoms may include extreme drowsiness, feeling light-headed, fainting, or coma.

## What should I avoid while taking Tranxene T-Tab (clorazepate)?

Do not drink alcohol while taking clorazepate. This medication can increase the effects of alcohol.

Clorazepate can cause side effects that may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be awake and alert.

Tell your doctor if you regularly use other medicines that make you sleepy (such as cold or allergy medicine, narcotic pain medicine, sleeping pills, muscle relaxers, and medicine for seizures, depression, or anxiety). They can add to sleepiness caused by clorazepate.

# Tranxene T-Tab (clorazepate) side effects

Get emergency medical help if you have any of these signs of an allergic reaction: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have any of these serious side effects:

- · confusion, unusual thoughts or behavior;
- · depressed mood, thoughts of suicide or hurting yourself;
- · muscle twitching, tremor; or
- problems with urination.

Less serious side effects may include:

- drowsiness, tiredness;
- · amnesia or forgetfulness;
- · dizziness;
- blurred vision;
- · feeling nervous or irritable;
- sleep problems (insomnia);
- muscle weakness, lack of balance or coordination;
- skin rash;
- nausea, vomiting, stomach pain, constipation; or
- · dry mouth; or
- · headache.

This is not a complete list of side effects and others may occur. Tell your doctor about any unusual or bothersome side effect. You may report side effects to FDA at 1-800-FDA-1088.

See also: Side effects (in more detail)



# What other drugs will affect Tranxene T-Tab (clorazepate)?

Before taking clorazepate, tell your doctor if you are taking any other anti-anxiety medications, or if you are using any of the following drugs:

- a barbiturate such as amobarbital (Amytal), butabarbital (Butisol), mephobarbital (Mebaral), secobarbital (Seconal), or phenobarbital (Luminal, Solfoton);
- an MAO inhibitor such as isocarboxazid (Marplan), phenelzine (Nardil), rasagiline (Azilect), selegiline (Eldepryl, Emsam), or tranylcypromine (Parnate);
- medicines to treat psychiatric disorders, such as chlorpromazine (Thorazine), haloperidol (Haldol), mesoridazine (Serentil), pimozide (Orap), or thioridazine (Mellaril);
- narcotic medications such as butorphanol (Stadol), codeine, hydrocodone (Lortab, Vicodin), levorphanol (Levo-Dromoran), meperidine (Demerol), methadone (Dolophine, Methadose), morphine (Kadian, MS Contin, Oramorph), naloxone (Narcan), oxycodone (OxyContin), propoxyphene (Darvon, Darvocet); or
- antidepressants such as amitriptyline (Elavil, Etrafon), amoxapine (Asendin), citalopram (Celexa), clomipramine (Anafranil), desipramine (Norpramin), doxepin (Sinequan), escitalopram (Lexapro), fluoxetine (Prozac, Sarafem), fluvoxamine (Luvox), imipramine (Janimine, Tofranil), nortriptyline (Pamelor), paroxetine (Paxil), protriptyline (Vivactil), sertraline (Zoloft), or trimipramine (Surmontil).

This list is not complete and there may be other drugs that can interact with clorazepate. Tell your doctor about all the prescription and over-the-counter medications you use. This includes vitamins, minerals, herbal products, and drugs prescribed by other doctors. Do not start using a new medication without telling your doctor.

Next  $\rightarrow$  Side Effects











### ▲ Add to My Med List

# More about Tranxene T-Tab (clorazepate)

- Side Effects
- During Pregnancy or Breastfeeding
- Dosage Information
- Drug Images
- Drug Interactions
- Support Group
- En Espanol
- 0 Reviews Add your own review/rating
- Consumer resources
- Tranxene T-Tab
- Tranxene T-Tab (Advanced Reading)
- Professional resources
- Tranxene T-Tab (FDA)
- Clorazepate Dipotassium (AHFS Monograph)
- Other formulations
- Tranxene
- Tranxene SD
- Related treatment guides
- Anxiety
- Alcohol Withdrawal
- Seizure Prevention

# Where can I get more information?

• Your pharmacist can provide more information about clorazepate.

!

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for

the indication prescribed.

Disclaimer: Every effort has been made to ensure that the information provided by Cerner Multum, Inc. ('Multum') is accurate, up-to-date, and complete, but no guarantee is made to that effect. Drug information contained herein may be time sensitive. Multum information has been compiled for use by healthcare practitioners and consumers in the United States and therefore Multum does not warrant that uses outside of the United States are appropriate, unless specifically indicated otherwise. Multum's drug information does not endorse drugs, diagnose patients or recommend therapy. Multum's drug information is an informational resource designed to assist licensed healthcare practitioners in caring for their patients and/or to serve consumers viewing this service as a supplement to, and not a substitute for, the expertise, skill, knowledge and judgment of healthcare practitioners. The absence of a warning for a given drug or drug combination in no way should be construed to indicate that the drug or drug combination is safe, effective or appropriate for any given patient. Multum does not assume any responsibility for any aspect of healthcare administered with the aid of information Multum provides. The information contained herein is not intended to cover all possible uses, directions, precautions, warnings, drug interactions, allergic reactions, or adverse effects. If you have questions about the drugs you are taking, check with your doctor, nurse or pharmacist.

ADUERTISEMENT

Copyright 1996-2012 Cerner Multum, Inc. Version: 5.04. Revision Date: 2010-12-15, 5:01:39 PM.

#### **DRUG STATUS**



Availability
Prescription only



Pregnancy Category Not classified



**CSA Schedule**Some potential for abuse

**Drug Class** 

Benzodiazepines

#### **Related Drugs**

Anxiety

Xanax, Lexapro, Cymbalta, alprazolam, Iorazepam, atenolol, Ativan, escitalopram, More...

**Alcohol Withdrawal** 

atenolol, diazepam, Valium, Librium, Tenormin, chlordiazepoxide, clorazepate, oxazepam, More...

**Seizure Prevention** 

 ${\bf clonazepam,\,Klonopin,\,diazepam,\,Valium,\,Lamictal,\,Topamax,\,lamotrigine,\,topiramate,\,More...}$ 

**CONSUMER UPDATES** 

Fighting Allergy Season with Medications

Medications for High Blood Pressure

Irritable Bowel Syndrome Treatments Aren't One-Size-Fits-All

Fighting Diabetes' Deadly Impact on Minorities

#### **Tranxene T-Tab Rating**

No Reviews - Be the first!

#### Tranxene T-Tab Images



Tranxene T-Tab (clorazepate) 7.5 mg





View all 6 images

#### **Related News and Articles**

Kids Who Aren't Ready for Kindergarten May Suffer Long-Term Consequences March 16, 2016

Anxiety, Depression May Reduce Women's Success With IVF: Study

March 15, 2016

**Could Too Much Cellphone Time Signal Anxiety, Depression?** 

March 15, 2016

#### **Related Questions & Answers**

Can you take both klonopin with tranxene t-tab for anxiety/panic disorder?

Clorazepate - I find that Tranxene 15mg has been taken off the market according to Recordati the new Has anyone been able to get brand Tranxene 15 mg since late 2014?

Read more questions... >





# Drugs.com Mobile Apps

The easiest way to lookup drug information, identify pills, check interactions and set up your own personal medication records. Available for Android and iOS devices.

#### **Explore Apps**



#### **SUPPORT**

**Help Center** 

**Frequent Questions** 

Sitemap

**Contact Us** 

#### **ABOUT**

**About Drugs.com** 

**Advertising Policy** 

**Content Submissions** 

**Drugs.com Blog** 

### **TERMS & PRIVACY**

**Editorial Policy** 

**Privacy Policy** 

Terms of Use

Attribution & Citations

#### CONNECT












Subscribe to receive email notifications whenever new articles are published.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex® (updated Feb 24th, 2016), Cerner Multum™ (updated Mar 1st, 2016), Wolters Kluwer™ (updated Mar 1st, 2016) and others.

To view content sources and attributions, please refer to our editorial policy.







We comply with the HONcode standard for trustworthy health information - verify here

Copyright © 2000-2016 Drugs.com. All rights reserved.